

İSTANBUL ANALYSIS SEMINARS

COMPACTNESS OF THE $\bar{\partial}$ -NEUMANN OPERATOR

Friedrich HASLINGER

Universität Wien
Faculty of Mathematics

Abstract: We consider the $\bar{\partial}$ -Neumann operator

$$N : L^2_{(0,q)}(\Omega) \rightarrow L^2_{(0,q)}(\Omega),$$

where $\Omega \subset \mathbb{C}^n$ is a bounded pseudoconvex domain, and

$$N_\varphi : L^2_{(0,q)}(\Omega, e^{-\varphi}) \rightarrow L^2_{(0,q)}(\Omega, e^{-\varphi}),$$

where $\Omega \subseteq \mathbb{C}^n$ is a pseudoconvex domain and φ is a plurisubharmonic weight function. N is the inverse to the complex Laplacian $\square = \bar{\partial}\bar{\partial}^* + \bar{\partial}^*\bar{\partial}$.

Using a general description of precompact subsets in L^2 -spaces we obtain a characterization of compactness of the $\bar{\partial}$ -Neumann operator, which can be applied to related questions about Schrödinger operators with magnetic field and Pauli and Dirac operators and to the complex Witten Laplacian. In this connection it is important to know whether the Fock space

$$\mathcal{A}^2(\mathbb{C}^n, e^{-\varphi}) = \{f : \mathbb{C}^n \rightarrow \mathbb{C} \text{ entire} : \int_{\mathbb{C}^n} |f|^2 e^{-\varphi} d\lambda < \infty\}$$

is infinite-dimensional, which depends on the behavior at infinity of the eigenvalues of the Levi matrix of the weight function φ .

In addition we discuss obstructions to compactness of the $\bar{\partial}$ -Neumann operator, and we describe, in some special cases, the spectrum of the \square -operator.

References

- [1] F. Haslinger, “Compactness for the $\bar{\partial}$ -Neumann problem: a functional analysis approach,” *Collect. Math.* **62** (2011), no. 2, 121-129.
- [2] F. Haslinger, “Spectrum of the $\bar{\partial}$ -Neumann Laplacian on the Fock space,” *J. Math. Anal. Appl.* **402** (2013), no. 2, 739-744.

Date: November 22, 2013

Time: 15:40

Place: Sabancı University, Karaköy Communication Center
Bankalar Caddesi 2, Karaköy 34420, İstanbul